



Muscle stimulation for the prevention of venous stasis during surgery

Kernohan, W. G., BROWN, JG., WARD, PE., & Mollan, RAB. (1989). Muscle stimulation for the prevention of venous stasis during surgery: (Conference abstract 1988; Book chapter 1989). In J. P. Paul, JC. Barbenel, JM. Courtney, & RM. Kenedi (Eds.), *Progress in Bioengineering: Proceedings of an international seminar held on the occasion of the 25th anniversary of the Strathclyde Bioengineering Unit, Glasgow in September 1988* (1 ed., Vol. 7, pp. 253-255). (Strathclyde bioengineering seminars). Adam Hilger.

[Link to publication record in Ulster University Research Portal](#)

Published in:
Progress in Bioengineering

Publication Status:
Published (in print/issue): 01/01/1989

Document Version
Publisher's PDF, also known as Version of record

General rights
Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.

MUSCLE STIMULATION FOR THE PREVENTION OF VENOUS STASIS DURING SURGERY

PE WARD, D.Phil, JG BROWN, F.R.C.S., WG KERNOHAN, Ph.D.,

RAB MOLLAN, M.D., F.R.C.S., F.R.C.S.I.

THE QUEEN'S UNIVERSITY OF BELFAST,

DEPARTMENT OF ORTHOPAEDIC SURGERY,

MUSGRAVE PARK HOSPITAL, STOCKMAN'S LANE, BELFAST BT9. 7JB.

Deep vein thrombosis is a major problem in the immediate post-operative phase following orthopaedic surgery, especially surgery involving the lower limbs. The venous flow rate is reduced while the patient is anaesthetised and this leads to venous stasis, which has been identified as an important factor involved in the formation of a thrombus.

In order that this might be prevented, a portable calf stimulator has been developed. This unit is easily applied and totally self contained. Two electrodes are attached to a non-conductive plastic case which contains the solid state circuitry and a rechargeable battery. The unit delivers a short pulse at a rate of 12/min. with the voltage variable from 70-100 volts. In position for use, the upper electrode lies distal to the popliteal fossa with the lower electrode over the insertion of the Achilles tendon. Such stimulation of the calf muscles causes them to contract helping push the blood up the deep calf veins towards the heart, thus preventing venous stasis.